Small Flock Management

Craig D. Coufal, Ph.D.

Associate Professor and Extension Specialist

Department of Poultry Science

ccoufal@poultry.tamu.edu





Getting Started in Poultry

- 1. Gather accurate information (see resources)
- 2. Find reputable source of birds
- 3. Plan well
 - Be prepared for all possibilities
 - Manure management/disposal
 - Dead bird disposal
 - Housing ventilation and weather extremes
 - Predator control
- 4. If for business, know the applicable rules and regulations

1. Informational Resources

- University/Extension
 - posc.tamu.edu
 - extension.org
 - Small and Backyard Flocks
 - Publications, webinars, Ask an Expert
- Government agencies
 - Texas Department of Agriculture (TDA), DSHS, USDA, CDC, etc.

2. Where do I get birds?


Reputable sources

- National Poultry Improvement Plan (NPIP) certified hatcheries (can order through US mail)
- Reputable breeders
 - Ask for documentation of disease testing (PT test)
- Not recommended:
 - Flea markets or "guy on the side of the road"
 - Ads in the paper
 - Friend of a friend

Mail Order Hatcheries

- Ideal Poultry Cameron, Texas
- Murray McMurray Iowa
- Morris Hatchery Miami, Florida
- Strombergs Minnesota
- Meyer Hatchery Ohio
- Welp Hatchery Iowa
- many more on internet


www.idealpoultry.com




What kind of bird?

- Depends on:
 - Type of production
 - Eggs
 - Meat
 - Both
 - Production expectations
 - Make a profit
 - Reduce food bill (not likely)
 - Personal use and satisfaction
 - What you like

White Egg Layers








LEGHORN — SINGLE COMB WHITE FEMALE

White Leghorn Production




Age in Weeks

White Egg Layers



Slide courtesy of Ideal Poultry

Brown Egg Layers



Silver Laced Wyandottes


New Hampshire Reds

Sex Links (not pictured)

Slide courtesy of Ideal Poultry

North Carolina Layer Test

Figure 30. Heritage Barred Plymouth Rock, Bi-weekly Percent Egg Production and Period Feed Consumption¹ in Hens kept on Range




1 kg per 100 Hens




Brown Leghorn-Cross




Hy-Line Brown Hen-Day Performance Graph



Green Egg Layers





ARAUCANAS SILVER DUCKWING — MALE

ARAUCANAS SILVER DUCKWING — FEMALE



Buff Orpingtons

Dual Purpose Breeds

Black Australorps



Dark Cornish






Delawares


Slide courtesy of Ideal Poultry

Meat Breeds





Black Broiler **Cornish Cross**



Red Broiler

Slide courtesy of Ideal Poultry

Disease Management

- Minimize contact between your birds and:
 - Manure source of bacteria
 - The ground source of parasites (worms)
 - Wild birds and rodents sources of anything bad
 - Eliminate habitat and attractants
 - Wild bird feeders and bird baths
 - Waterfowl ponds
 - Clean up spilled feed to reduce attraction
 - Pets source of bacteria and pests (especially fleas)

Introducing new birds to a flock

- Quarantine from existing flock for at least 3 weeks
 - Not in same building or pen
 - As far away as possible
 - Monitor for disease symptoms, particularly respiratory
 - Handle sick or quarantined birds AFTER the other birds.
 - Wash hands thoroughly and disinfect boots
 - Not recommended to comingle birds of significantly different ages

Vaccination

- Minimum recommended vaccinations:
 - Marek's disease
 - Fowl pox
- Other possibilities:
 - Newcastle disease
 - Infectious bursal disease (IBD) (gumboro)
- Vaccines can be ordered on the internet or can be administered at the hatchery





Fowl pox



Medications

- Antibiotics do not treat viral infectins
- Almost all antibiotics are NOT approved by FDA for laying hens
 - Prior to January 1, 2017 OTC antibiotics approved for hens producing eggs for food included:
 - Bacitracin zinc
 - Bacitracin methylene disalicylate (BMD)
 - Chlortetracycline
 - Tylosin

Veterinary Feed Directive

- As of January 1, 2017, all <u>medically important</u> <u>antibiotics</u> to be used in feed or water for food animal species require a Veterinary Feed Directive (VFD) or a prescription.
- FDA requires veterinary oversight whenever medically important antibiotics are administered to any food animal species via feed or water, **even if the animals are not intended for food production.** From pet rabbits and pigs, to backyard poultry, to large livestock farms, the same restrictions apply.
- The only antibiotics considered not important are Bacitracin, Meccadox, Tiamulin (Denegard), Narasin, and Bambermycin.

Drugs Transitioning From OTC to VFD Status (from FDA website)

Established drug name	Examples of proprietary drug name(s)
chlortetracycline (CTC)	Aureomycin, CLTC, CTC, Chloratet, Chlorachel, ChlorMax, Chlortetracycline, Deracin, Inchlor, Pennchlor, Pfichlor
chlortetracycline/sulfamethazine	Aureo S, Aureomix S, Pennchlor S
chlortetracycline/sulfamethazine/ penicillin	Aureomix 500, Chlorachel/Pficlor SP, Pennchlor SP, ChlorMax SP
hygromycin B	Hygromix
lincomycin	Lincomix
oxytetracycline (OTC)	TM, OXTC, Oxytetracycline, Pennox, Terramycin
oxytetracycline/neomycin	Neo-Oxy, Neo-Terramycin
penicillin	Penicillin, Penicillin G Procaine
sulfadimethoxine/ormetoprim	Rofenaid, Romet
tylosin	Tylan, Tylosin, Tylovet
tylosin/sulfamethazine	Tylan Sulfa G, Tylan Plus Sulfa G, Tylosin Plus Sulfamethazine
virginiamycin	Stafac, Virginiamycin, V-Max

Drugs Transitioning From OTC to VFD Status (from FDA website)

Established drug name	Examples of proprietary drug name(s)
chlortetracycline (CTC)	Aureomycin, CLTC, CTC, Chloratet, Chlorachel, ChlorMax, Chlortetracycline, Deracin, Inchlor, Pennchlor, Pfichlor
chlortetracycline/sulfamethazine	Aureo S, Aureomix S, Pennchlor S
chlortetracycline/sulfamethazine/ penicillin	Aureomix 500, Chlorachel/Pficlor SP, Pennchlor SP, ChlorMax SP
hygromycin B	Hygromix
lincomycin	Lincomix
oxytetracycline (OTC)	TM, OXTC, Oxytetracycline, Pennox, Terramycin
oxytetracycline/neomycin	Neo-Oxy, Neo-Terramycin
penicillin	Penicillin, Penicillin G Procaine
sulfadimethoxine/ormetoprim	Rofenaid, Romet
tylosin	Tylan, Tylosin, Tylovet
tylosin/sulfamethazine	Tylan Sulfa G, Tylan Plus Sulfa G, Tylosin Plus Sulfamethazine
virginiamycin	Stafac, Virginiamycin, V-Max

Medications

- Almost all antibiotics NOT approved by FDA for laying hens
 - Approved for hens producing eggs for food : bacitracin zinc, BMD, <u>chlortetracycline</u>, and tylosin Veterinarian prescription only
- NO wormers were approved for laying hens prior to 2018 (
- Fenbendazole (sold as Safe-guard®) now approved as of January, 2018 to treat intestinal worms

Disease Diagnosis

- Please do not call the Poultry Science
 Department for an over-the-phone diagnosis
 - We can handle all other aspects of poultry production (nutrition, housing, fertility, incubation, brooding, etc.)
- Who to call?
 - 1. Local veterinarian
 - 2. Texas A&M Veterinary Medical Diagnostic Laboratory (TVMDL)
 - 3. Reportable disease Texas Animal Health Commission (TAHC)

TVMDL

- tvmdl.tamu.edu
 - 3 poultry diagnostic labs in Texas
 - College Station 979-845-3414
 - * Gonzales 830-672-2834
 - Center 936-598-4451
 - Can preform necropsy for a fee
 - See website for more information

One More Big Concern Biosecurity and Disease





Avian influenza (AI)

- Virus is carried by wild birds
- 2 forms:
 - Low pathogenic (LPAI)
 - Highly pathogenic (HPAI)
 - Highly contagious to domestic poultry
 - May cause high mortality in poultry flocks
- Spring, 2015 outbreak of HPAI across Midwest US
 - Largest in US history
 - Devastating economic impacts
 - Consumers impacted by higher egg prices

Prevention of AI

- What can we do?
 - Vaccination not yet an option

• Only option: must prevent infection

- How?
 - Sound biosecurity is the only tool we have

What is Biosecurity?

- Set of management practices designed to prevent the introduction of disease
 - "Protection from infection"
 - http://healthybirds.aphis.usda.gov/





Good Biosecurity

- 3 main components:
 - Isolation prevent contact with infected vectors (wild birds and animals, rodents, pets, etc.)
 - **Traffic control** limit access to flock to reduce exposure risk (people and vehicles)
 - Sanitation disinfect anything that comes into contact with your flock (vehicles, equipment, coops, footwear, clothing, etc.)



Biosecurity Suggestions

- Avoid visiting more than one flock per day
- Shower, change clothes and disinfect shoes between visits
- Use shoe covers to keep shoes clean
- Use dedicated clothes and shoes not worn anywhere else for checking on flock

Biosecurity Suggestions

- Discourage presence of wild birds
- Do not keep waterfowl and chicken/turkeys on the same premises
- Do not share equipment, coops or old feed
- Consider all-in, all-out style of management

Virulent Newcastle Disease (VND)

- Formerly known as Exotic Newcastle Disease (END)
- Virus
- Vaccination recommended
- Biosecurity essential
- Endemic to Mexico
- Current outbreak in southern California has spread to Utah (381 confirmed cases since May, 2018)

Mycoplasmosis

- Cause *Mycoplasma gallisepticum, M. synoviae*
 - "Bacteria" with no cell wall (prokaryote)
- Disease
 - Respiratory rales, coughing, nasal discharge, conjunctivitis, sinusitis
 - Secondary bacterial infections (colibacillosis)
- Treatment
 - Antibiotic injections or water administration
 - Tylosin, enrofloxacin, erythromycin, lincomycin-spectinomycin
- Prevention
 - Live and inactivated vaccines

Internal parasites

Common intestinal parasites of chickens Roundworms Tapeworms Threadworms Cecal worms

• Treat with fenbendazole (Safe-guard)

Housing



Functions of Housing

- Protection from weather extremes
 - precipitation
 - winter and summer
- Protection from predators
 particularly at night
- For laying hens, a place to nest and lay eggs
- For chicks, proper brooding environment for early growth period
- Maximize efficiency



Better, but can be improved




Good Design with Adjustable Doors




Location considerations

- Adequate drainage a must
- Proximity to habitat for wild birds and predators
- Harborage for pests and rodents
 - eliminate trash and brush piles, old equipment
- Natural air flow (breeze)
- Visibility from your house
- Lack of visibility from your neighbor's house









Principles of Housing Design

- Shape of house
 - Rectangular superior to square
 - Natural ventilation
 - want to minimize the distance air must travel through naturally ventilated houses





- Mechanical ventilation
 - holding ventilation rate constant, air speed increases as cross-sectional area decreases



Principles of Housing Design

- Orientation
 - Long axis east-west
 - Minimizes wall area directly facing sun
 - For naturally ventilated houses:
 - minimizes direct sunlight shining into house
 - takes advantage of prevailing south wind in summer

Principles of Housing Design

Materials

- Metal is fine for exterior walls and roof
- Metal not recommended for interior surfaces
 - metal is an excellent conductor of heat (radiant heat transfer)
 - Insulation can help
- Cleaning and disinfection should be considered for interior surfaces
 - lower porosity materials are easier to clean
 - paint can help to seal wood surfaces for easier cleaning

But what about mobile coops or "chicken tractors"?



Poultry: Science and Practice, 1951

196

Poultry: Science and Practice



Fig. 104. A well-constructed range shelter built on skids to facilitate moving.



Highly recommend anchoring

Things to consider with mobile coops

- May have to move daily, depending on density of birds
 - Enough land??
- Heat/Cold stress
 - Air flow
 - Shade
 - Moisture
- Predators
- Rain and ground moisture
- Still must brood baby chicks in a building

Nipple drinker systems are suitable for all chickens (not turkeys)

- automated
 - reduced time and labor
- more sanitary than trough or bell drinkers
- Iow pressure
- less spillage
 = drier manure/ litter




Laying hens on nipple drinkers



Water Equipment Options

 FarmTek catalog (also online)



Simple Nipple Drinkers with a Bucket





Creating your own Poultry Nipple Bucket Drinker is as easy as 1-2-3!

- Drill holes in the bottom of the sealed bucket with an 11/32" drill bit (purchased locally). We suggest three holes, spaced accordingly, for maximum efficiency.
- VVhen installing nipples, place rubber grommet in drilled hole first, followed by the actual nipple. Grommet is included with each nipple. Wetting the nipple first eases the application.
- Hanging or fastening your drinker will depend on the location and application.
 We recommend using rope, cable or chain. All are sold separately. Ask your National Account Manager for more details.

PVC Pipe style



